

HIV RISK ASSESSMENT

<u>Risk Factors</u>	<u>Yes</u>	<u>No</u>
I. Since 1978, have you		
a. - used IV drugs and shared needles/syringes?	_____	_____
b. - immigrated from a country with a high prevalence of HIV infection (Haiti or Central African countries)?	_____	_____
c. - engaged in prostitution	_____	_____
d. - received blood/blood products prior to 4/85?	_____	_____
e. - been raped?	_____	_____
f. - had evidence of HIV infection?	_____	_____
g. - had genital ulcer disease?	_____	_____
(For Women Only)		
h. - been artificially inseminated prior to 11/86?	_____	_____
(For Men Only)		
I. - received clotting factor for treatment of hemophilia prior to 4/85?	_____	_____
II. Since 1978, have you been a sex partner of		
a. - an IV drug user?	_____	_____
b. - a person who received blood/blood products prior to 4/85?	_____	_____
c. - an entrant or immigrant from a country with a prevalence of HIV infection (Haiti or Central African countries)?	_____	_____
d. - a person who has AIDS, ARC or HIV infection?	_____	_____
e. - a prostitute?	_____	_____
(For Men Only)		
f. - another man?	_____	_____
(For Women Only)		
g. - a man who has had sex with another man?	_____	_____
III. Have you had more than one sex partner in the last six months?		
	_____	_____

A "yes" answer to any of the above questions puts the patient at increased risk

The following non-specific signs and symptoms may indicate HIV related disease:

- * Change in mental status or motor functions that interferes with daily activities
- * Swollen glands
- * Weight loss 10% of baseline body weight not due to diet or exercise
- * Repeated night sweats and/or fever for more than 30 days in absence of another illness
- * Diarrhea for more than 30 days in absence of another illness
- * Weakness
- * Dry cough/shortness of breath
- * Purplish-blue spots on skin